

On Simplest Hamiltonian Systems

Pavol Ševera

*Dept. of Theoretical Physics, Charles University
V Holešovičkách 2 182 00 Prague, Czech Republic*

Abstract

Simple Hamiltonian systems, such as mathematical pendulum or Euler equations for rigid body, are solved without computation. It is nothing but a joke but maybe you will find it nice.

There are some simple mechanical systems, such as mathematical pendulum or Euler equations for rigid body, admitting solution in elliptic functions. These functions are one of the most beautiful things in elementary complex analysis, especially when they are treated in Riemannian way, using their global properties and singularities and avoiding computation. As a simple, but maybe charming example, we present a geometrical solution of the afore-mentioned systems.

Let our phase space be an irreducible quadratic surface $F = 0$ in a 3-dimensional affine space A_3 . The symplectic form is given as follows: choose a trivector ϵ_3 ; then $\sigma = \langle \epsilon_3, dF \rangle$ (when restricted to $F = 0$) is the bivector field on the phase space giving the Poisson structure (i.e. the inverse of the symplectic form). It is invariant with respect to unimodular affine transformations preserving F and, if all singularities of $F = 0$ are hidden at infinity, this property fixes σ up to a constant multiple. If H is the Hamiltonian, then

$$v = \langle \epsilon_3, dF \wedge dH \rangle \quad (1)$$

is the evolution field.

Suppose that H is a time-independent quadratic function on A_3 . The best known examples are Euler equations (the phase space a sphere) and (possibly rotating) mathematical pendulum (the phase space a cylinder). Any energy level, say $H = 0$, is a spatial quartic curve. As soon as it is regular, it is an elliptic curve (if we project it from its arbitrary point, we obtain a regular planar cubic). We will see that (1) is regular on the curve, including the complex points and the points at infinity. Thus the evolution is nothing but addition on the elliptic curve and the system is solved in elliptic functions.

We will restate (1) using projective geometry. Let V be a vector space, C a 2-dimensional cone in V and c the corresponding curve in the projective space $\mathbf{P}(V)$. A vector field v on c can be realized in V as a bivector field b on C , homogenous of degree 2. The correspondence is as follows:

Let $P \in C$ and let $p \in c$ be the corresponding point in $\mathbf{P}(V)$ and let v' be a vector at P projected to v at p (v' is unique up to addition of a constant multiple of P). Then $b = P \wedge v'$. Also, if $\alpha \in V^*$ is such that $\alpha(P) = 1$ then $\langle b, \alpha \rangle$ is a possible choice of v' .

Now let A_3 be realized in a 4-dimensional V by an equation $\alpha = 1$. Also let f, h be the homogenizations of F, H . Let ϵ_4 be the unique tetravector on V such that $\epsilon_3 = \langle \epsilon_4, \alpha \rangle$. Then, according to (1), on the cone $f = h = 0$,

$$b = \langle \epsilon_4, df \wedge dh \rangle. \quad (2)$$

Equation (2) is a priori valid on $\alpha = 1$, but in fact it is valid everywhere on C , as $df \wedge dh$ has the right homogeneity.

We see that b , and thus v , is everywhere finite.

If $df \wedge dh = 0$ anywhere on the curve, the situation is even more favourable. The curve is necessarily singular (on an elliptic curve, v is nowhere zero) and thus it is a rational curve or a reducible curve composed of rational ones. All the following assertions are obtained by considering the curve as a limiting case of a regular one.

If the curve is irreducible with a cusp, equations of motion are solved in rational functions: if the cusp is identified with ∞ , the evolution is simply addition.

If the curve has a simple double point, solution can be given in exponentials: if the coalescing points are identified with 0 and ∞ , the evolution is multiplication. The exponential is real or imaginary according to the location of the double point with respect to the other real points.

If the curve decomposes (to a line and a twisted cubic, or to two (possibly reducible) conics), the evolution is rational or exponential according to the location of points of intersection. If a component is double, $df \wedge dh = 0$ identically there and the evolution is trivial.

References

- [1] V.I. Arnold: *Mathematical Methods of Classical Mechanics*, Springer, Berlin-Heidelberg-New York 1978.
- [2] F. Klein: *Vorlesungen über die Entwicklung der Mathematik in 19. Jahrhundert*, Springer, Berlin 1926.